



# Addicted to plastic: Microplastic pollution and prevention



This briefing outlines global issues relating to microplastic pollution and its prevention. Microplastics are particles that are smaller than 5mm and are formed by the fragmentation of larger plastic items or are intentionally or unintentionally released in the form of manufactured beads, granules, fibres and fragments. Once in the environment they are very difficult to remove and have the potential to accumulate in soil, freshwater and marine environments causing a range of known and unknown impacts.

## Contents

Summary .....	3
Recommendations.....	5
Our addiction to plastic.....	6
In detail .....	7
Sources of microplastics.....	7
Impacts of microplastics.....	8
Removal of microplastics through water treatment processes.....	10
Accumulation of microplastics.....	11
Initiatives to prevent pollution.....	11
Acknowledgements.....	14
References.....	14

CIWEM is the leading independent  
Chartered professional body for water and  
environmental professionals, promoting  
excellence within the sector

### **CIWEM**

106-109 Saffron Hill

London EC1N8QS

Charity registration No 1043409 (England  
and Wales) SC038212 Scotland

[www.ciwem.org](http://www.ciwem.org)

Cover image: Trudie Ballantyne,  
*Straws*. CIWEM's Environmental  
photographer of the Year  
competition

## Summary

### **The majority of microplastics are unintentionally lost to the environment**

There are seven main sources of microplastics (tyres, synthetic materials, marine coatings, road markings, personal care products, city dust and losses during plastic production). Most are unintentionally lost through abrasion, weathering or unintended spills, or originate from the mismanagement of waste containing plastics. The largest proportion result from the washing of synthetic textiles and from the abrasion of road markings and tyres from driving.

### **More than half of microplastic losses will remain on land and in soils**

Whilst the accumulation of plastics in the marine environment has received worldwide attention, it is estimated that more than half of microplastic losses will remain on land. Removal at wastewater treatment works has proven to be highly effective, however, due to the large volumes of wastewater that are processed, millions of microplastics are still being released back into the freshwater environment each day. The large quantities that are retained at the treatment works may then be transferred back into the environment via the spreading of sewage sludge on land for agriculture.

### **Microplastics are likely to have environmental impacts but data is sparse**

Studies in the marine environment have shown microplastics can be ingested by marine animals leading to physical harm and reproductive or toxic effects. The adverse environmental effects of the polymers and additives which make up the microplastics following dietary uptake are not fully known.

Although most research has focused on marine environments, freshwater systems may be at greater risk from primary microplastic contamination due to their closer proximity to wastewater treatment plants and plastic processing factories. Yet the biological effects of microplastics in freshwater species is reasonably sparse.

The possible negative environmental impact associated with the application of sewage sludge containing microplastics to farmland requires much greater research: Microplastics could provide a medium for exotic species and pathogens to grow on or release plasticisers and POPs (persistent organic pollutants), which thereby have the potential to enter the human food chain via crops or livestock grazing.

### **A global issue**

Initiatives to tackle plastic and microplastic pollution fall far short of what is needed. An international protocol or regulatory framework to guide co-operation between governments, plastics manufacturers, the waste management and water industries, consumers and companies using plastics in their products is necessary to solve issues around design, manufacturing, use, reuse and disposal of plastics.



### Prevention is the best, and most likely the cheapest, solution to reduce microplastic pollution

Urgent measures are needed to address microplastic pollution at source, as once released into the environment there are few, if any, practicable means by which these pollutants can be removed. Banning manufacturers from including primary microplastics such as microbeads in 'wash off' personal care products such as face scrubs, toothpastes and shower gels is welcome, however these plastics are a minor contributor to the overall problem (<4 per cent). As other losses are unintentional, from product use and maintenance, through abrasion, weathering or unintended spills, preventing them will be all the more difficult. The plastics industry must do more to prevent losses in production via Operation Clean Sweep and the textile industry and white goods manufacturers should act to reduce the amount of fibres ending up in wastewater.

### A new plastics' strategy is needed - designing for reuse

Outside of the European Union the UK will need to address plastics with its own strategy. Plastics can easily form part of a circular economy with reuse or recycling if they are designed appropriately and are properly recovered and managed when they reach the end of their life. Methods should include improved product design and substitution, extended producer responsibility and deposit return schemes.



*Courtesy Belat Rand: Searching in garbage, CIWEM's Environmental Photographer of the Year*



## Recommendations

1. **Governments** to set high standards and improve recovery and recycling of plastics to minimise the quantity of secondary microplastics that reach the environment.
2. **Research** is needed into the fate and transport mechanisms of microplastics within the environment and into which polymers are the most damaging and under what conditions, so that these can be most effectively addressed.
3. **Governments** to provide incentives (financial or otherwise) for the use of alternatives that are shown to be less damaging and tackle the issue of single-use packaging items like plastic film which can be neither reused nor recycled.
4. **Industry** to develop potentially commercially viable plastics and plastic alternatives that are less damaging to the environment, particularly for tyres, road markings and synthetic fabrics.
5. **Local authorities** to review road markings and highway drainage to prevent microplastics entering the freshwater environment.
6. **Regulators** to enforce existing restrictions on the use of hazardous additives and polymer ingredients under the EU REACH regulations, encourage producers to use more benign alternatives where possible, and in the longer term take a more sector-based approach to the assessment and regulation of chemicals related to plastics.
7. **Governments** to promote good practice in plastic use and disposal and increase public awareness of plastic use (wider than personal care items e.g. littering, washing synthetic clothing, windblown litter), reuse and recycling capabilities.
8. **Developed countries** to target development assistance on waste management to alleviate what in many parts of the world is an immense problem.



*Courtesy David Cheresna: Globalisation  
CIWEM's Environmental Photographer of  
the Year*

## Our addiction to plastic

Society's adoption of plastics as a substitute for traditional materials has expanded extremely rapidly since large-scale production began in the 1950s. Each year over 322 million tonnes of plastic are produced globally and this is increasing<sup>1</sup>.

Plastics are an incredibly versatile material with many uses for which alternatives are not viable, either functionally or financially. They are made from a group of large molecules called polymers which vary in characteristics such as buoyancy, toxicity and degradability. Plastics may also contain additives designed to change the properties of the end product.

Durability is a common feature of most plastics, and it is this property, combined with an unwillingness or inability to manage end-of-life plastic effectively that has resulted in waste plastics and microplastics becoming a global problem. Many plastic items are so cheap they can be thrown away without financial consequence to the consumer: only 9 per cent of the global plastic waste produced since the 1950s has been recycled with 79 per cent in landfills or in the wider environment<sup>2</sup>.

The plastic waste problem is growing rapidly in developing countries too, partly due to increased affordability of products made from plastic and partly due to vastly inadequate infrastructure to manage the waste problem. Over three billion people do not have access to waste collection and disposal facilities.

Microplastics are formed by the fragmentation and weathering of larger plastic items or are in the form of manufactured beads, granules, fibres and fragments.


Each year billions of particles of microplastic find their way into the environment and being hard to degrade, they accumulate. They have become ubiquitous and are abundant not only in all of the five ocean subtropical gyres but also in Arctic Sea-ice, deep sea sediments and uninhabited Pacific islands.

## DEFINITIONS

**Microplastics** consist of particles smaller than 5mm right down to microscopic particles. Those that are 1-5mm are usually referred to as large microplastics and <1mm are small microplastics. They are categorised as primary or secondary:

**Primary microplastics** include industrial scrubbers used in blast cleaning, plastic powders used in moulding, plastic nanoparticles used in a variety of industrial processes, micro-beads in cosmetic formulations and soaps and in detergents for washing machines.

**Secondary microplastics** are formed by the fragmentation and weathering of larger plastic items during the use of products such as textiles, paint and tyres, or once these or other plastic items (bags, bottles etc.) have been released into the environment.



## In detail

### Sources of microplastics

The International Union for Conservation of Nature and Natural Resources (IUCN) describe seven major sources of microplastics: tyres, synthetic materials, marine coatings, road markings, personal care products, city dust and plastic pellets (incidents during the manufacturing, transport and recycling of plastics). Apart from personal care products, most microplastics are unintentionally lost through abrasion, weathering or unintended spills. Secondary microplastics largely originate from the mismanagement of waste during the disposal of products containing plastics.

Two thirds of primary microplastics originate from road runoff (tyres, road markings and pellets incidents on land) (figures 1 and 2). Another important source is from the washing of clothes made from synthetic materials. Research shows the washing of manmade fabrics can potentially release over 700,000 fibres from an average wash load of 6kg<sup>3</sup>.




Figure 1. Global releases to the world oceans. IUCN<sup>4</sup>. Contribution of different pathways to the release of microplastics

Microplastics once in the environment can be transported via wind, commercial and domestic discharges to a sewer, runoff into rivers, runoff into combined sewer systems and runoff directly into lakes and oceans. The IUCN estimate that of the microplastic losses, 48 per cent end up in the ocean and around 52 per cent are trapped in soils when wastewater sewage sludge is used as fertiliser and/or when particulates are washed from the road pavement.

Their presence on land and in rivers, lakes, groundwater and the ocean is influenced by a combination of environmental factors including exposure to UV radiation, buoyancy and by the properties of the polymer from which they are



made. The properties of microplastics in the oceans differ substantially from those in sewage and run off as there has generally been more opportunity for the breakdown of larger plastic fragments into secondary microplastics.



*Figure 2. High friction plastics are used for junctions, cycle lanes and on roads with high traffic volumes*

## Impacts of microplastics

In the ocean many microplastic particles are found suspended in water, where they can enter the food chain by being ingested by filter feeders, ranging from zooplankton to baleen whales, or ingested by benthic species where denser particles sink to the bottom.

Of the few available studies, a study on anchovies has suggested that plastics are mistaken for food due to the chemical signature plastic debris acquires in the ocean<sup>5</sup>, another that polystyrene nanoparticles have severe effects on both behaviour and metabolism in fish<sup>6</sup>. There are some studies demonstrating chemical adsorption to microplastics and desorption in the gut following ingestion, based on chemical analyses and also whole organism effects<sup>7</sup>.

Although most research has focused on marine environments, freshwater systems may be at greater risk from primary microplastic contamination, due to them being closer to point sources (such as wastewater treatment plants and plastic processing factories). Data on the biological effects of microplastics in freshwater species is reasonably sparse. The accumulation of other freshwater contaminants on microplastics' surfaces is of special interest because ingestion has the potential to increase the chemical exposure.



Organisms that ingest microplastics particles lack a digestive system that can degrade them. Fibres can clump and knot blocking the digestive tract of small organisms in a similar way that larger plastics do in larger organisms. Also the nutrition of small organisms is potentially undermined if they mistake plastics for food. The adverse environmental effects of the polymers and additives which make up the microplastics following dietary uptake are not fully known. Although additives will have been tested as a requirement under the EU REACH regulation, these will have been aquatic ecotoxicity studies with waterborne exposure, and not dietary studies.

The potential for toxic or hormonal disruptive effects of microplastics are attributed by UNEP to persistent organic pollutants (POPs) and other persistent, bioaccumulative and toxic chemicals that may have been used as additives in plastic manufacture, and to the ability of microplastic particles to attract and concentrate harmful such organic pollutants with which they come into contact. Microplastics could also provide a medium for exotic species and pathogens to grow on, for example microorganisms developing biofilms on microplastics particles.



## Removal of microplastics through water treatment processes

### Drinking water

The process of turning raw water into drinking water involves a three stage treatment which includes coagulation, flocculation and filtration. This is primarily to remove cryptosporidium spores ( $5\mu\text{m}$ ) from drinking water and so water treatment plants should therefore remove microplastics down to this size and prevent them going into drinking water. Treatment would not generally remove particles below this size so there could be implications for nanoplastics entering drinking water ( $<100\text{nm}$ ).

### Wastewater

Removal at wastewater treatment works is highly dependent on the density and size of the particle<sup>8</sup>. Microplastics that are dense enough to settle are periodically removed from settling tanks along with other settled solids. Those that float are scraped from the surface of settling tanks before being mixed with the settled solids. Microplastics that remain in suspension may escape further treatment filters, such as those used in tertiary treatment plants, and are therefore discharged from the works into surface water or rivers as part of effluent discharge.

Wastewater treatment works are not specifically designed to remove microplastics. However removal by conventional primary and secondary wastewater treatment technologies have shown to be very effective overall, (studies show around 97-99 per cent are removed)<sup>9</sup>. Yet despite this large reduction, due to the large volumes of wastewater processed each day, a large





treatment works could still release approximately 900,000 to 3600 million microplastics per day to freshwater and marine environments<sup>10</sup>.

As treatment can leave 99 per cent of microplastics in sludge, this is a potentially large source of pollution. Microplastic retention in sewage sludge and subsequent application of sewage sludge to terrestrial systems for agricultural reasons (sludge can supply a large part of the nitrogen or phosphorus needed by most crops) may lead to the transfer of microplastics and/or chemicals to soil used in growing food<sup>11</sup>. EU legislation requires sludge to be treated to protect against health hazards, for example by lime stabilisation, anaerobic digestion, composting, or thermal drying, but there is limited evidence of these being able to remove microplastics<sup>12</sup> and there is currently no specific regulation for microplastics. Anaerobic digestion should be further investigated as a remediation technique.

## Accumulation of microplastics

The majority of microplastics research to date has focussed on the marine environment; in 2017 a survey of 279 UK shorelines found 73 per cent contained plastic pellets<sup>13</sup>. Yet each year it is believed that between 473,000 and 910,000 metric tonnes of plastic waste is released and retained within land-based environments. This equates to between 4 and 23 times the amount estimated to be deposited in oceans<sup>14</sup>. The only way they may be naturally removed is by wind or run-off so once incorporated beneath the surface into the soil there may be limited chance of this happening. They may eventually, in part be returned to the aquatic environment<sup>15</sup>.

Microplastics that are removed in wastewater treatment plants become concentrated in sewage sludge. The possible negative environmental impact associated with the application of sewage sludge containing microplastics to farmland has not been extensively researched. However microplastics degrade slowly and as a result can accumulate in the soil. Both plasticisers and POPs (persistent organic pollutants) could be released from plastics, and thereby enter the human food chain via crops or livestock grazing. POPs have been shown to be taken up by plants, but at a lower rate than heavy metals, for example.

## Initiatives to prevent pollution

### International

A responsibility to protect the marine environment is enshrined in the UN Convention on the Law of the Sea which requires states to individually or jointly as appropriate prevent, reduce and control pollution of the marine environment from any source. The UN Sustainable Development Goals also include a target to prevent and significantly reduce marine pollution of all kinds, including marine

litter. However most Member States are failing to take responsibility for primary and secondary plastic pollution.

In 2017 voluntary commitments were made by a number of countries whose rivers transport plastic waste to the ocean: Thailand pledged to implement proper waste disposal and encourage environmentally-friendly alternatives to plastic packaging as part of its recent 20-year pollution management strategy and Indonesia pledged to reduce plastic waste by 70 per cent by 2025.

## Europe

The European Marine Strategy Framework Directive addresses the issue of marine litter, including plastics. Microplastics are covered by Descriptor 10 of Commission Decision 2010/477/EU, which defines the good environmental status of marine waters<sup>16</sup>. It recommends that ideally microplastics are reduced at origin. In contrast, the Water Framework Directive applying to European inland waters does not specifically refer to plastic litter. As microplastics are an emerging issue, they may be considered within the Water Framework Directive's list of Priority Substances or under the Hazardous Substances Directive in the future which would have significant implications for the water industry. A number of substances are controlled under EU chemicals policy (REACH) which restricts some plastics from certain uses.

A Directive to reduce the use of light-weight plastic carrier bags was adopted in 2015 to reduce the consumption of and to phase out bags that fragment rather than degrade<sup>17</sup>. Plastics are now one of the five priority areas addressed in the EU action plan for the Circular Economy. Plastic waste already needs to be collected separately, but the Package proposes raising the recycling target for plastic packaging to 55 per cent, and reducing landfilling to no more than 10 per cent by 2030. It will set out a strategy to assess the challenges across the entire life cycle of plastics, improve recycling, cut marine litter, and remove potentially dangerous chemicals towards the end of 2017.

## Microbeads

Some effort is being made to reduce the occurrence of microbeads added to 'wash off' personal care products such as face scrubs, toothpastes and shower gels. A joint statement to the European Commission by Sweden, Belgium, The Netherlands, Austria and Luxembourg, urged a ban on microplastics in cosmetics and detergents. Bans are under consideration in individual EU Member States, including Ireland and France. The UK has committed to introducing a microbead ban in cosmetics and personal care products starting in 2018<sup>18</sup>. Although a minor contributor to the overall problem (estimates range from 0.01 to 4.1 per cent of plastics in the marine environment<sup>19</sup>) this is an easy win as the microplastics are easily be replaced with far less damaging alternatives. At the time of writing some large cosmetic manufacturers had removed microbeads from their personal care products ahead of imminent bans. Measures should also be taken to avoid



redefinition of the plastic beads used in personal care products so that they are able to continue under another guise.

## UK

Charging for single-use plastic bags to reduce consumption has been highly successful in the UK with more than nine million fewer plastic bags used (around 83 per cent reduction) since the government introduced a 5p charge and around 40 per cent fewer have been found on beaches<sup>20</sup>. The government should now consider plastic bottle deposit return schemes which have shown to have extremely high recovery rates (~90 per cent) in Germany, Sweden and Norway.

Outside of the European Union the UK will need to address plastics with its own strategy. Following the publication of the Litter Strategy for England the government will follow with a new national anti-littering campaign in 2018, working with industry and the voluntary sector to drive behaviour change. The UK will need to take a far more proactive approach to waste policy, delivering the circular economy cannot be left to the market alone.



## Acknowledgements

Lead author Laura Grant, Senior Policy Adviser. Produced with thanks to members from CIWEM's Water Supply and Quality Panel, Wastewater and Biosolids Panel, Contaminated Land Network and Waste Management Panel. Thanks to Dr Alice Horton, CEH and Nic Thompson for input and advice.

[policy@ciwem.org](mailto:policy@ciwem.org)

August 2017

*Briefings represent the Institution's views on issues at a particular point in time. It is accepted that situations change as research provides new evidence. It should be understood, therefore, that CIWEM briefings are under constant review and that previously held views may alter and lead to revisions. Briefings are produced as a consensus report and do not represent the view of individual members of CIWEM.*

## References

- 1 European Commission. 2017. Plastics factsheet
- 2 Geyer, R, Jambeck, J and Lavender Law, K. 2017. Production, use, and fate of all plastics ever made. Science Advances Vol 3, No. 7
- 3 Napper, I.E. and Thompson, R.C. 2016. Release of synthetic microplastic plastic fibres from domestic washing machines: Effects of fabric type and washing conditions. Marine Pollution Bulletin, 112, 39-45
- 4 Boucher, J., Friot, D. 2017. Primary microplastics in the oceans: a global evaluation of sources. International Union for Conservation of Nature
- 5 Matthew S. Savoca, Chris W. Tyson, Michael McGill, Christina J. Slager. 2017. Odours from marine plastic debris induce food search behaviours in a forage fish. Proc. R. Soc. B 2017
- 6 Mattsson K1 ,Ekvall MT, Hansson LA, Linse S, Malmendal A, Cedervall T. 2015. Altered behavior, physiology, and metabolism in fish exposed to polystyrene nanoparticles. Environ Sci Technol. 6;49(1):553-61.
- 7 Browne M.A., Niven S.J., Galloway T.S., Rowland S.J., Thompson R.C. 2013. Microplastic moves pollutants and additives to worms, reducing functions linked to health and biodiversity. Curr Biol 23:2388-2392, and Rochman C.M., Hoh E., Kurobe T., Teh S.J. 2013. Ingested plastic transfers hazardous chemicals to fish and induces hepatic stress. Sci Rep 3:3263, and Teuten E.L., Saquing J.M., Knappe D.R.U., Barlaz M.A., Jonsson S., Bjorn A. 2009. Transport and release of chemicals from plastics to the environment and to wildlife. Philosophical Transactions of the Royal Society B 364:2027-2045, and Bakir A., Rowland S.J., Thompson R.C. 2014. Enhanced desorption of persistent organic pollutants from microplastics under simulated physiological conditions. Environ Pollut 185:16-23.
- 8 Carr, S. A.; Liu, J.; Tesoro, A. G. 2016. Transport and fate of microplastic particles in wastewater treatment plants. Water Res. 91, 174–182
- 9 Magnusson K & Norén F IVL Swedish Environmental Research Institute: Screening of microplastic particles in and down-stream a wastewater treatment plant, 2014, Number C55, and Murphy, F., Ewins, C., Carbonnier, F., Quinn, B. 2016. Wastewater Treatment Works (WwTW) as a Source of Microplastics in the Aquatic Environment. Environmental Science & Technology 50, and Talvitie, J., Mikola, A., Setälä, O., Heinonen, M., Koistinen, A. 2017. How well is microliter purified from wastewater? -A detailed study on the stepwise removal of microliter in a



- 
- tertiary level wastewater treatment plant. *Water Research* 109, 164-172, and Mintenig, S.M., Int-Veen, I., Loder, M.G., Primpke, S., Gerds, G. 2017. Identification of microplastic in effluents of waste water treatment plants using focal plane array-based micro-Fourier-transform infrared imaging. *Water Research* 108, 365-372
- 10 Horton, A. 2017. Microplastics in freshwater systems – presence, origins and ecological effects Centre for Ecology and Hydrology. Presentation to CIWEM Wastewater and Biosolids panel and Ziajahromi, S., Neale, P. A., & Leusch, F. D. 2016. Wastewater treatment plant effluent as a source of microplastics: review of the fate, chemical interactions and potential risks to aquatic organisms. *Water Science and Technology*, 74 (10), 2253-2269.
  - 11 Eerkes-Medrano, D. et al, 2015. Microplastics in freshwater systems: A review of the emerging threats, identification of knowledge gaps and prioritisation of research needs. *Water Research*
  - 12 Mahon, A.M., O'Connell, B., Healy, M.G. et al. 2017. Microplastics in Sewage Sludge: Effects of Treatment. *Environmental Science & Technology*, 51 (2): 810–818
  - 13 Nurdle free oceans <http://www.nurdlehunt.org.uk/take-part/nurdle-map.html>
  - 14 Horton, A., et al. 2017. Microplastics in freshwater and terrestrial environments: evaluating the current understanding to identify the knowledge gaps and future research needs. *Science of the total environment* 586, 127-141
  - 15 Nizzetto, L., Futter, M., Langaas, S. 2016. Are agricultural soils dumps for microplastic of urban origin? *Environmental Science and Technology*
  - 16 Galgani F, Hanke G, Werner S, De Vrees L. 2013. Marine litter within the European Marine Strategy Framework Directive. *ICES Journal of Marine Science* 70: 1055–1064
  - 17 Directive (EU) 2015/720 of the European Parliament and of the Council of 29 April 2015 amending Directive 94/62/EC as regards reducing the consumption of lightweight plastic carrier bag
  - 18 Defra. 03/09/2016 Microbead ban announced to protect sea life and Environment Secretary pledges action on ocean plastics. Defra press release 21 July 2017
  - 19 Defra. 2016. Evidence to the EAC on Environmental impacts of microplastics
  - 20 Environment Secretary pledges action on ocean plastics. Defra press release 21 July 2017 and Marine Conservation Society. 2016. Great British Beach Clean.